

# LONG ABDOMINAL AORTIC STENOSIS – A CASE OF TAKAYASU ARTERITIS


David Campos-Correia<sup>1\*</sup>, Ana Coutinho Santos<sup>1</sup>, Vítor Bettencourt<sup>2</sup>, Carla Saraiva<sup>1</sup>

<sup>1</sup>Radiology Department, Centro Hospitalar de Lisboa Ocidental

<sup>2</sup>Vascular Surgery Department, Centro Hospitalar de Lisboa Ocidental

\*Contacto Autor: davidcorreia7@gmail.com

A 77-year-old female Caucasian patient with known Takayasu's arteritis diagnosed at 20 years of age was admitted to the emergency department due to diffuse sudden-onset abdominal pain. On physical examination, femoral pulses were feeble. Laboratory results were unremarkable. Abdominal CT angiography showed a long abdominal predominantly infra-renal aortic stenosis (Figures 1 and 2).



**Figure 1**

Multislice computed tomography angiography. Thin maximum intensity projection (MIP) axial images. **A** – markedly reduced lumen of the abdominal infra-renal aorta, with a maximum diameter of less than 1cm, with circumferential extensive calcifications. **B** – At a lower level, marked luminal reduction with peripheral and circumferential hypodense thickening of the aortic wall can be seen in keeping with the diagnosis of Takayasu Arteritis.



**Figure 2**

**A** – 2D curved reformatted thin MIP coronal image shows collateral circulation and the inferior mesenteric artery with a larger calibre than usual. **B** – 3D volume-rendered (VR) reformatted image shows the long stenosis extending from the infra-renal aorta to the iliac arteries with extensive calcifications. The large calibre of the inferior mesenteric artery and aortic arch involvement can also be appreciated.